



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Recurrence of myopericarditis following mRNA COVID-19 vaccination in a male adolescent

Tomohiko C. Umei, MD, Yoshikazu Kishino, MD, PhD, Yasuyuki Shiraishi, MD, PhD, Taku Inohara, MD, PhD, Shinsuke Yuasa, MD, PhD, Keiichi Fukuda, MD, PhD



Journal of the  
Journal de la  
Canadian  
Cardiovascular  
Society  
Société  
canadienne  
de cardiologie

PII: S2589-790X(21)00351-6

DOI: <https://doi.org/10.1016/j.cjco.2021.12.002>

Reference: CJCO 471

To appear in: *CJC Open*

Received Date: 15 November 2021

Accepted Date: 5 December 2021

Please cite this article as: T.C. Umei, Y. Kishino, Y. Shiraishi, T. Inohara, S. Yuasa, K. Fukuda, Recurrence of myopericarditis following mRNA COVID-19 vaccination in a male adolescent, *CJC Open* (2022), doi: <https://doi.org/10.1016/j.cjco.2021.12.002>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Inc. on behalf of the Canadian Cardiovascular Society.

*Case Reports*

**Recurrence of myopericarditis following mRNA COVID-19 vaccination in a male adolescent**

**Short title:** Myocarditis recurrence after COVID-19 vaccine

**Authors:**

Tomohiko C. Umei, MD, Yoshikazu Kishino\*, MD, PhD, Yasuyuki Shiraishi, MD, PhD,  
Taku Inohara, MD, PhD, Shinsuke Yuasa, MD, PhD, Keiichi Fukuda, MD, PhD

**Institution:**

Department of Cardiology, Keio University School of Medicine

**\*Corresponding Author:**

Yoshikazu Kishino, MD, PhD

Department of Cardiology, Keio University School of Medicine

35 Shinanomachi Shinjuku-ku, Tokyo, Japan, 160-8582

E-mail: ykishino@z7.keio.jp TEL: +81-3-3353-1211 FAX: +81-3-5843-6167

**Word count:** 879 words (from the beginning of the Introduction to the end of the Discussion )

**Keywords:** myopericarditis, vaccine, COVID-19

## 1    **Brief Summary**

2    We present a case of recurrent myopericarditis in a young man following an initial episode  
3    of influenza virus-induced myopericarditis after receiving the mRNA-1273 COVID-19  
4    vaccine. This case highlights the importance of assessing patients' backgrounds, which  
5    may help elucidate the mechanism and risk factors of vaccine-induced myopericarditis.

6



## Abstract

Coronavirus disease 2019 (COVID-19) is caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). It has spread worldwide, resulting in health and economic crises. Vaccination against SARS-CoV-2 infection is considered a valid prevention measure to control this pandemic. There have been reports of cases of myopericarditis following mRNA COVID-19 vaccination. We present a case of a 20-year-old man with recurrent myopericarditis following an initial episode of influenza virus-induced myopericarditis and after receiving a second dose of the mRNA-1273 Moderna COVID-19 vaccine. Careful attention should be paid to patients with a history of myocarditis following COVID-19 vaccination.

## 1 Introduction

2 Coronavirus disease 2019 (COVID-19) is caused by severe acute respiratory syndrome  
3 coronavirus 2 (SARS-CoV-2) that has spread worldwide, resulting in health and  
4 economic crises. Vaccination against SARS-CoV-2 is considered a protective approach  
5 with more than 95% efficacy to control the COVID-19 pandemic <sup>1</sup>. However, some cases  
6 of myopericarditis have recently been reported as complications of mRNA COVID-19  
7 vaccination with an incidence of 1 case per 10,000–100,000 inoculations <sup>2</sup>. Although  
8 these events are rare, cases were much more common following a second dose, especially  
9 in young male adolescents <sup>2</sup>. Herein, we present a case of a 20-year-old male adolescent  
10 with recurrent acute myopericarditis five years after an initial episode of acute influenza  
11 virus-induced myopericarditis and two days after receiving a second dose of the mRNA-  
12 1273 Moderna COVID-19 vaccine.

## 14 Case

15 A 20-year-old male adolescent had a history of seasonal influenza virus-induced  
16 myopericarditis in 2016 (data not shown). He presented to our hospital in July 2021 with  
17 chief complaints of fever and severe sharp chest pain in deep breathing improved by  
18 sitting up. The patient had been well and free from any flu-like illnesses a couple of  
19 months before admission. Two days before admission, he received a second dose of the  
20 mRNA-1273 Moderna COVID-19 vaccine. Initial vital signs were normal, except for a  
21 high body temperature of 38.8°C. Electrocardiography (ECG) showed global ST–T wave  
22 elevation (**Figure 1A**). Transthoracic echocardiography revealed regional hypokinesis of

the anteroseptal wall, an ejection fraction of 53.8%, and no pericardial effusion (**Supplemental Movie 1A**). Serum creatinine kinase (CK) and troponin T levels were elevated. Polymerase chain reaction testing for SARS-CoV-2 and antigen tests for influenza virus were negative. Paired serum samples were negative for viruses that were possible causes of myocarditis, including coxsackievirus, adenovirus, and human parvovirus B19. Accordingly, the patient received conservative management, including administration of oral loxoprofen at 160 mg/day for one week. Serum CK (1190 IU/L) and troponin T (0.710 ng/mL) levels peaked on day 2 after admission. Cardiac magnetic resonance imaging (cMR) showed no apparent findings on T1 and T2 mapping. However, high signal intensity on T2-weighted in the inferolateral wall in regions with late gadolinium enhancement was suggestive of acute myocarditis (**Figure 2A and 2B**). Changes in ECG findings revealed a pseudonormalized ST-T pattern followed by T-wave inversion in precordial leads (**Figure 1B and 1C**). His symptoms gradually resolved, and he was discharged on day 5. He remained stable and his left ventricular function improved one month after discharge (**Supplemental Movie 1B**).

## Discussion

Vaccination against SARS-CoV-2 is considered a potent approach to control the COVID-19 pandemic. The efficacy of the mRNA COVID-19 vaccine has been reported to be 95%<sup>1</sup>. However, some cases of myopericarditis have recently been reported as complications of mRNA COVID-19 vaccination with an incidence of 1 case per 10,000–100,000 inoculations<sup>2</sup>. Although these events are as rare as extrapulmonary manifestations of influenza, cases were much more common following a second dose of mRNA COVID-

19 vaccination, especially in young male adolescents <sup>2</sup>.

Historically, vaccine-induced myopericarditis was reported as rare adverse events, such as after receiving smallpox or influenza virus vaccinations <sup>3</sup>. Circulating heart-reactive autoantibodies have been reported in a high percentage of patients with myocarditis and have been linked to its pathogenesis. Another possible mechanism for vaccine-induced myocarditis might have been molecular similarities between anti-virus antibodies and self-antigens. The mRNA-1273 vaccine is a lipid nanoparticle-encapsulated mRNA-based vaccine that encodes the prefusion-stabilized full-length spike protein of SARS-CoV-2. Of note, antibodies against SARS-CoV-2 spike proteins have been experimentally shown to cross-react with structurally similar human peptide sequences, including alpha-myosin <sup>4</sup>. In addition, a history of infection can be related to the magnitude of immune response to the current infection in a phenomenon referred to as “original antigenic sin”. A case of repeated pericarditis after vaccination for influenza has been reported <sup>5</sup>. This concept refers to cross-reacting immunity due to past infections with similar virus strains, which must be considered when interpreting immune responses to infections and vaccinations. However, there have been some reports that common viruses including influenza virus were poor sources of cross-reactive immunity to SARS-CoV-2 <sup>6</sup>. Since we did not perform testing for viral genomes or autoantibodies in heart tissues, a direct relationship between influenza virus-induced and COVID-19 mRNA vaccine-induced myocarditis cannot be definitively established. In the setting of stable uncomplicated myocarditis, cMR rather than endomyocardial biopsy is recommended. However, biopsy would be further considered for elucidating the mechanism <sup>7</sup>.

COVID-19 has spread worldwide and vaccination against COVID-19 is an effective

approach to prevent infection and aggravation. There have been some reports of myocarditis induced by COVID-19 mRNA vaccine. However, the exact mechanism of mRNA vaccine-induced myocarditis has yet to be determined. Our case was a patient with recurrent myocarditis following an initial episode of influenza infection and after receiving the mRNA-1273 Moderna COVID-19 vaccine. It is not that COVID-19 vaccination is not recommended for patients with prior myocarditis. However, careful watch would be considered. Accumulation of myopericarditis cases following mRNA vaccination, and assessing their backgrounds, particularly regarding the immune system, may help elucidate the mechanism and risk factors for developing this adverse event.

### **Novel teaching points**

□ Both influenza myocarditis and COVID-19 mRNA vaccine-induced myocarditis are rare but might occur consecutively.

□ Careful watch is important after COVID-19 vaccination for cases with a history of myocarditis.

### **Conclusion**

We experienced a case of a 20-year-old man with recurrent myopericarditis following an initial episode of influenza virus-induced myopericarditis and after receiving a

second dose of the mRNA-1273 Moderna COVID-19 vaccine. Careful watch should be considered after COVID-19 vaccination for cases with a history of myocarditis.

#### Acknowledgements

None.

#### Funding

None.

#### Disclosures

None.

#### References

1. Polack FP, Thomas SJ, Kitchin N, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med* 2020;383:2603-15.
2. Luk A, Clarke B, Dahdah N, et al. Myocarditis and Pericarditis following COVID-19 mRNA Vaccination: Practical Considerations for Care Providers. *Can J Cardiol* 2021.
3. Su JR, McNeil MM, Welsh KJ, et al. Myopericarditis after vaccination, Vaccine Adverse Event Reporting System (VAERS), 1990-2018. *Vaccine* 2021;39:839-45.
4. Vojdani A, Kharrazian D. Potential antigenic cross-reactivity between SARS-CoV-2 and human tissue with a possible link to an increase in autoimmune diseases. *Clin Immunol* 2020;217:108480.
5. Mei R, Raschi E, Poluzzi E, Diemberger I, De Ponti F. Recurrence of pericarditis after influenza vaccination: a case report and review of the literature. *BMC Pharmacol Toxicol* 2018;19:20.
6. Reche PA. Potential Cross-Reactive Immunity to SARS-CoV-2 From Common Human Pathogens and Vaccines. *Front Immunol* 2020;11:586984.

- 1 7. Tschöpe C, Ammirati E, Bozkurt B, et al. Myocarditis and inflammatory  
2 cardiomyopathy: current evidence and future directions. Nat Rev Cardiol  
3 2021;18:169-93.  
4

## Figure Legends

### Figure 1. ECG changes during hospitalization and after discharge.

ECG obtained upon admission showing global ST-T wave elevation (A). Changes in ECG reveals an improved ST-T elevation on day 2 (B), followed by T-wave inversion in precordial leads one month after discharge (C). *ECG, electrocardiogram.*

### Figure 2. cMR imaging on day 3.

cMR imaging shows regional hyperintensity on T2-weighted (A) and linear sub-epicardial LGE (B) in the inferolateral wall. *cMR, cardiac magnetic resonance; LGE, late gadolinium enhancement.*

### Supplemental Video 1. 2D speckle tracking echocardiography.

TTE performed on admission day shows hypokinesis of the antero-septal wall with no pericardial effusion (A). TTE findings one month after discharge shows an almost normal left ventricular function (B). *2D, 2-dimensional; TTE, transthoracic echocardiography.*





